0790

PB 204 147

COMPUTER MAPPING
as an aid in
AIR POLLUTION STUDIES

VOLUME ONE: SUMMARY REPORT

APPENDIX: COMPUTER GRAPHICS

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
Springfield, Va. 22151

A

157

BIBLIOGRAPHIC DATA SHEET	1. Report No. APTD-0790	2.	3. Recipient's Accession No.
4. Title and Subtitle Computer Mapping as an aid in Air Pollution Studies Volume One: Summary Report - Appendix: Computer Graphics			5. Report Date April 1970 6.
7. Author(s) John C. Goodrich, Editor and Gail Howrigan, Associate Editor			8. Performing Organization Rept. No.
9. Performing Organization Name and Address Graduate School of Design Harvard University Cambridge, Massachusetts 02138			10. Project/Task/Work Unit No. 11. Contract/Grant No. 68A-2405D
12. Sponsoring Organization Name and Address National Air Pollution Control Administration U. S. Public Health Service U. S. Department of Health, Education, and Welfare Washington, D. C. 20201			13. Type of Report & Period Covered 14.

15. Supplementary Notes DISCLAIMER - This report was furnished to the Office of Air Programs by Graduate School of Design, Harvard University, Cambridge, Massachusetts 02138 in fulfillment of Grant No. 68A-2405D

16. Abstracts

An appendix which contains a complete set of all graphics shown in volume I and volume II of a study of computer mapping as applied to air pollution studies.

17. Key Words and Document Analysis. 17a. Descriptors

Air pollution
Computer programs
Mapping
Graphic methods

17b. Identifiers/Open-Ended Terms

17c. COSATI Field/Group 13B

18. Availability Statement Unlimited	19. Security Class (This Report) UNCLASSIFIED	21. No. of Pages 50
	20. Security Class (This Page UNCLASSIFIED	22. Price

LABORATORY FOR COMPUTER GRAPHICS
AND SPATIAL ANALYSIS

Graduate School of Design
Harvard University
Cambridge, Massachusetts

This report was funded by:

Demonstration Grant 68A-2405D
National Air Pollution Control Administration
U. S. Public Health Service
U. S. Department of Health, Education, and Welfare

John C. Goodrich, Editor
Gail Howrigan, Associate Editor

Rosalie Alexander
Carol Chase
Candace Howes
Whitmore John
Linda Peacock

April 1970

COMPUTER MAPPING

as an aid in

AIR POLLUTION STUDIES

Personnel

Principal Investigators

Howard T. Fisher	Project Director (Associate Director of The Laboratory for Computer Graphics and Spatial Analysis)
John C. Goodrich	Associate Project Director
William Warntz	(Director of The Laboratory for Computer Graphics and Spatial Analysis)
Allan H. Schmidt	(Associate Director of The Laboratory for Computer Graphics and Spatial Analysis)

Case Study Investigators

Norma Bogen	St. Louis Project
Paul Jack Dangermond	California and Puget Sound Studies
Steven Fosburg	Statistical Investigation
Jean Maurice Granger	Montreal Project
Donald Shepard	St. Louis Project/Program Investigation
Elia Werczberger	Kansas City Project

Programming Support and Development

Chang-I Hua
Lannon Leiman
Kathleen Reine

Research Assistants

Whitmore John
Johnny Kirkland
Helen Mansfield
Adrian Thomas

Ellen Bussard
Robert Cartwright
Katherine Mickey
Warnie Webster

Administrative and Secretarial Support

Malcolm Fitzpatrick
Gail Howrigan
Linda Peacock
Laurence Yont

Regina McDonald
Donna L. Moorhead
Claudia S. Rosen
Judith Strong



FIGURE A-1

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Census Tract Centers



FIGURE A-2

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Measurement Stations — Winter Network — 1963-64
a. Except Sulfur Dioxide b. Sulfur Dioxide



FIGURE A-3

ST. LOUIS REGION STUDY

St. Louis City — Land Use Maps

a. Percent Land Area in Residential Use

b. Percent Land Area in Public and Semipublic Use

c. Percent of Land Area in Commercial Use



FIGURE A-4

ST. LOUIS REGION STUDY
St. Louis City -- Land Use Maps
a. Percent Land Area in Industrial Use
b. Percent Land Area in Streets and Alleys
c. Percent Land Area Vacant



FIGURE A-5

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
24-hour SO_2 Readings — 19-Station Network
a. Arithmetic Means, Winter 1963-64
b. Arithmetic Means, Winter 1964-65



FIGURE A-6

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Differences between A-5a. and b.





FIGURE A-7

ST. LOUIS REGION STUDY

St. Louis Interstate Air Pollution Study Area

24-hour SO_2 Readings — 40-Station Network

a. Arithmetic Means — Winter 1964-65

b. Geometric Means — Winter 1964-65









FIGURE A-8

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Suspended Particulates by High-Volume Air Sampler
Fall 1963 (Weekday Schedule)
a. Minimum b. Maximum









FIGURE A-9

ST. LOUIS REGION STUDY

St. Louis Interstate Air Pollution Study Area
Suspended Particulates by High-Volume Air Sampler
Fall 1963 (Weekday Schedule)

a. Arithmetic Mean b. Geometric Mean



C SUSPENDED PARTICULATES BY HIGH-VOLUME AIR SAMPLER
C WINTER 1963-1964 (WEEKDAY SCHEDULE)
C MINIMUM

DATA VALUE EXTREMES ARE 12.00 93.00

ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL
(*MAXIMUM* INCLUDED IN HIGHEST LEVEL ONLY)

	1	2	3	4	5
MINIMUM	12.00	25.50	47.70	59.50	71.30
MAXIMUM	24.40	35.70	46.90	71.30	93.00

PERCENTAGE OF TOTAL ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL

1	2	3	4	5
15.31	6.54	20.74	30.62	26.79

FREQUENCY DISTRIBUTION OF DATA POINT VALUES IN EACH LEVEL

LEVEL	1	2	3	4	5
SYMBOLS					
FREQ.	3	3	4	3	3

LABORATORY FOR COMPUTER GRAPHICS AND SPATIAL ANALYSIS
HARVARD UNIVERSITY CAMBRIDGE, MASSACHUSETTS

SOURCE - INTERSTATE AIR POLLUTION STUDY, PHASE 2, PROJECT REPORT VOLUME 2, AIR QUALITY MEASUREMENTS PAGES 25,26 U.S.DEPARTMENT OF HEALTH,EDUCATION, AND WELFARE-DEC.1966

OUTLINE OF MAP APPROXIMATES LIMITS OF INTERSTATE AIR POLLUTION STUDY AREA

GRID SYSTEM DEFINED ON RIGHT AND BOTTOM BORDER

MISSISSIPPI AND MISSOURI RIVERS INDICATED BY WHITE LINE



C SUSPENDED PARTICULATES BY HIGH-VOLUME AIR SAMPLER
C WINTER 1963-1964 (WEEKDAY SCHEDULE)
C MAXIMUM

DATA VALUE EXTREMES ARE 150.00 1337.00

ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL
(*MAXIMUM* INCLUDED IN HIGHEST LEVEL ONLY)

	1	2	3	4	5
MINIMUM	150.00	197.40	257.00	316.00	420.90
MAXIMUM	197.30	257.00	316.00	420.90	1337.00

PERCENTAGE OF TOTAL ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL

1	2	3	4	5
3.90	9.04	6.97	8.04	77.10

FREQUENCY DISTRIBUTION OF DATA POINT VALUES IN EACH LEVEL

LEVEL	1	2	3	4	5
SYMBOLS					
FREQ.	3	3	4	3	3

LABORATORY FOR COMPUTER GRAPHICS AND SPATIAL ANALYSIS
HARVARD UNIVERSITY CAMBRIDGE, MASSACHUSETTS

SOURCE - INTERSTATE AIR POLLUTION STUDY, PHASE 2, PROJECT REPORT VOLUME 2, AIR QUALITY MEASUREMENTS PAGES 25,26 U.S.DEPARTMENT OF HEALTH,EDUCATION, AND WELFARE-DEC.1966

OUTLINE OF MAP APPROXIMATES LIMITS OF INTERSTATE AIR POLLUTION STUDY AREA

GRID SYSTEM DEFINED ON RIGHT AND BOTTOM BORDER

MISSISSIPPI AND MISSOURI RIVERS INDICATED BY WHITE LINE

FIGURE A-10

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Suspended Particulates by High-Volume Air Sampler
Winter 1963-64 (Weekday Schedule)
a. Minimum b. Maximum



C SUSPENDED PARTICULATES BY HIGH-VOLUME AIR SAMPLER
C WINTER 1963-1964 (WEEKDAY SCHEDULE)
C ARITHMETIC MEAN

DATA VALUE EXTREMES ARE 57.28 204.32

ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL
("MAXIMUM" INCLUDED IN HIGHEST LEVEL ONLY)

MINIMUM					
	92.16	130.46	169.39	196.58	185.68

PERCENTAGE OF TOTAL ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL

8.29	10.74	18.40	16.02	46.55

FREQUENCY DISTRIBUTION OF DATA POINT VALUES IN EACH LEVEL

LEVEL	1	2	3	4	5
SYMBOLS	.	:	+	E	E
FREQ.	3	3	4	3	3

LABORATORY FOR COMPUTER GRAPHICS AND SPATIAL ANALYSIS
HARVARD UNIVERSITY CAMBRIDGE, MASSACHUSETTS

SOURCE - INTERSTATE AIR POLLUTION STUDY, PHASE 2, PROJECT REPORT
VOLUME I, AIR QUALITY MEASUREMENTS
PUBLIC HEALTH SERVICE PUBLICATION NO. 999-AP-28
U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE-DEC. 1966

OUTLINE OF MAP APPROXIMATES LIMITS OF INTERSTATE AIR POLLUTION STUDY AREA

GRID SYSTEM DEFINED ON RIGHT AND BOTTOM BORDER

MISSISSIPPI AND MISSOURI RIVERS INDICATED BY WHITE LINE



C SUSPENDED PARTICULATES BY HIGH-VOLUME AIR SAMPLER
C WINTER 1963-1964 (WEEKDAY SCHEDULE)
C GEOMETRIC MEAN

DATA VALUE EXTREMES ARE 52.16 206.57

ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL
("MAXIMUM" INCLUDED IN HIGHEST LEVEL ONLY)

MINIMUM					
	89.15	127.82	166.82	194.25	184.38

PERCENTAGE OF TOTAL ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL

8.74	16.11	25.90	25.63	23.62

FREQUENCY DISTRIBUTION OF DATA POINT VALUES IN EACH LEVEL

LEVEL	1	2	3	4	5
SYMBOLS	.	:	+	E	E
FREQ.	3	3	4	5	5

LABORATORY FOR COMPUTER GRAPHICS AND SPATIAL ANALYSIS
HARVARD UNIVERSITY CAMBRIDGE, MASSACHUSETTS

SOURCE - INTERSTATE AIR POLLUTION STUDY, PHASE 2, PROJECT REPORT
VOLUME I, AIR QUALITY MEASUREMENTS
PUBLIC HEALTH SERVICE PUBLICATION NO. 999-AP-28
U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE-DEC. 1966

OUTLINE OF MAP APPROXIMATES LIMITS OF INTERSTATE AIR POLLUTION STUDY AREA

GRID SYSTEM DEFINED ON RIGHT AND BOTTOM BORDER

MISSISSIPPI AND MISSOURI RIVERS INDICATED BY WHITE LINE

FIGURE A-11

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Suspended Particulates by High-Volume Air Sampler
Winter 1963-64 (Weekday Schedule)
a. Arithmetic Mean b. Geometric Mean





FIGURE A-12

ST. LOUIS REGION STUDY

St. Louis Interstate Air Pollution Study Area

Suspended Particulates by High-Volume Air Sampler

July 1963-June 1964 (Random Schedule)

a. Minimum b. Maximum





FIGURE A-13

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Suspended Particulates by High-Volume Air Sampler
July 1963-June 1964 (Random Schedule)
a. Arithmetic Mean b. Geometric Mean



FIGURE A-14

ST. LOUIS REGION STUDY

St. Louis Interstate Air Pollution Study Area

Comparison Map — Winter 1964-65

SO_2 Level versus Population Density

a. Using .03 Level b. Using .04 Level



FIGURE A-15

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Comparison Map
SO_2 Level versus Density of Population under Age Ten
a. Using .03 Level b. Using .04 Level



FIGURE A-16

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Comparison Map
SO_2 Level versus Density of Population over Age Sixty-five
a. Using .03 Level b. Using .04 Level



FIGURE A-17

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Comparison Map
a. SO_2 (Winter 1963-64) Using .04 Level, Suspended Particulates (July 1963-June 1964) Using 75 μ/m^3 Level, and Population Density
b. SO_2 (Winter 1963-64) Using .04 Level, Suspended Particulates (July 1963-June 1964) Using 120 μ/m^3 Level, and Population Density



FIGURE B-1

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Surface of 2-hour Sulfur Dioxide Readings
December 20, 1964 — 4, 6, and 8 a.m.



FIGURE B-2

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Surface of 2-hour Sulfur Dioxide Readings
December 20, 1964 — 10, 12 a.m. and 2 p.m.



FIGURE B-3

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
2-hour SO_2 Surface in Oblique View
December 20, 1964 — 4 and 6 a.m.



FIGURE B-4

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
2-hour SO_2 Surface in Oblique View
December 20, 1964 — 8 and 10 a.m.



FIGURE B-5

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
2-hour SO_2 Surface in Oblique View
December 20, 1964 — noon and 2 p.m.



FIGURE B-6

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Surface of 2-hour Sulfur Dioxide Readings
December 21, 1964 — 4, 6, and 8 a.m.



FIGURE B-7

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
Surface of 2-hour Sulfur Dioxide Readings
December 21, 1964 — 10, 12 a.m. and 2 p.m.





FIGURE B-8

ST. LOUIS REGION STUDY
St. Louis Interstate Air Pollution Study Area
a. Regression Estimate of 24-hour SO_2 Readings
b. 24-hour SO_2 Readings Corresponding to Above





FIGURE B-9

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Population Density
Contour Map and Oblique View



FIGURE B-10

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Population Density — Oblique Views
a. 90° Rotation b. 180° Rotation



FIGURE B-11

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Percent of Population under 10 Years of Age



FIGURE B-12

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Percent of Population over 65 Years of Age



FIGURE B-13

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Median School Years Completed



FIGURE B-14

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
a. Percent of Population Non-white
b. Percent of Families with Incomes less than $3000/yr.



FIGURE B-15

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Percent of Families with Incomes less than $3000/yr.
Oblique View





FIGURE B-16

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Density of Families with Incomes less than $3000/yr.





FIGURE B-17

ST. LOUIS REGION STUDY

Standard Metropolitan Statistical Area — 1960 Census

a. Percent of Families with Incomes from $3000 to $7999/yr.

b. Percent of Families with Incomes of $8000/yr. or More





FIGURE B-18

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Percent of Housing Units Deteriorating and/or Dilapidated
a. Contour View b. Oblique View



FIGURE B-19

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Percent of Housing Units with 1.01 or more Persons per Room



FIGURE B-20

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Percent of Housing with 1 Unit/Structure



FIGURE B-21

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
a. Percent of Housing with 2 Units/Structure
b. Percent of Housing with 3 or 4 Units/Structure



FIGURE B-22

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
a. Percent of Housing with 5 to 9 Units/Structure
b. Percent of Housing with 10 or more Units/Structure



FIGURE B-23

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Percent of Housing Built in 1939 or Earlier



FIGURE B-24

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Percent of Housing Built 1940-1949



FIGURE B-25

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
Percent of Housing Built 1950-1960





FIGURE B-26

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
a. Density of Population over 65 Years of Age
b. Oblique View of Same



FIGURE B-27

ST. LOUIS REGION STUDY
Standard Metropolitan Statistical Area — 1960 Census
a. Density of Population under 10 Years of Age
b. Oblique View of Same



FIGURE C-1

ST. LOUIS REGION STUDY
Power Plant Study Area
Measured SO_2 Concentrations — 22 January 1965



FIGURE C-2

ST. LOUIS REGION STUDY
Power Plant Study Area
Projected 1970 Population Density
Mapped by Census Tracts



POPULATION DENSITY, PROJECTED 1970, POPULATION PER SQUARE MILE
BASED ON 1960 CENSUS TRACTS AND 1970 COUNTY PROJECTIONS BY
THE EAST-WEST GATEWAY COORDINATING COUNCIL, ST LOUIS AREA

SOURCE - U. S. CENSUSES OF POPULATION AND HOUSING, 1960
CENTERS OF CENSUS TRACTS DEFINED ACCORDING TO U. S. CENSUS
DEFINITION IN TERMS OF LATITUDE AND LONGITUDE
MISSISSIPPI AND MISSOURI RIVERS INDICATED BY WHITE LINE

DATA VALUE EXTREMES ARE 17.62 10102.93

ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL
(*MAXIMUM* INCLUDED IN HIGHEST LEVEL ONLY)

MINIMUM	0.00	500.00	2000.00	5000.00	10000.00	20000.00
MAXIMUM	500.00	2000.00	5000.00	10000.00	20000.00	70000.00

PERCENTAGE OF TOTAL ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL

	0.71	2.14	4.29	7.14	14.29	71.43

FREQUENCY DISTRIBUTION OF DATA POINT VALUES IN EACH LEVEL

LEVEL	1	2	3	4	5	6
SYMBOLS						
FREQ.	420	72	60	66	10	8

FIGURE C-3

ST. LOUIS REGION STUDY
Power Plant Study Area
Projected 1970 Population Density
Mapped by Grid Coordinates



FIGURE C-4

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Moderate Winds from South
SO_2 Concentrations without Regard to "Social Cost"



SULFUR DIOXIDE CONCENTRATIONS FOR MINIMAL EXPOSURE -- ST LOUIS, FPWR
BASED ON OPTIMAL DISPATCHING OF ELECTRIC POWER STATIONS, 1970
USING GIFFUSEON MODEL WITH MODERATE WIND FROM THE SOUTH, NEUTRAL STAB.

DATA VALUE EXTREMES ARE 0.00 9.74

ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL
(MAXIMUM INCLUDED IN HIGHEST LEVEL ONLY)

MINIMUM	BELOW	0.09	0.17	0.25	0.92	1.58	6.26	ABOVE 9.00
MAXIMUM	0.01	0.09	0.17	0.25	0.92	1.58	6.26	9.00

PERCENTAGE OF TOTAL ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL

3.12	3.12	4.25	12.90	29.00	50.00	

FREQUENCY DISTRIBUTION OF DATA POINT VALUES IN EACH LEVEL

LEVEL	0	1	2	3	4	5	6	7
SYMBOLS	----------	----------	----------	----------	----------	----------	----------	----------
FREQ.	332	89	56	61	40	24	7	1

FIGURE C-5

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Moderate Winds from South
SO_2 Concentrations for Minimal Exposure



FIGURE C-6

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Moderate Winds from South
Improvement in SO_2 through "Social Cost" Dispatching



FIGURE C-7

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Moderate Winds from South
Density of Exposure to Pollution without Regard to "Social Cost"



FIGURE C-8

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Moderate Winds from South



FIGURE C-9

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Moderate Winds from South
Improvement in Population Exposure through "Social Cost" Dispatching



FIGURE C-10

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Light Winds from West
SO_2 Concentrations without Regard to "Social Cost"



FIGURE C-11

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Light Winds from West
SO_2 Concentrations without Regard to "Social Cost"



IMPROVEMENT IN AIR POLLUTION RESULTING FROM 'SOCIAL COST' DISPATCHING
SULFUR DIOXIDE CONCENTRATIONS IN PARTS PER HUNDRED MILLION
USING DIFFUSION MODEL WITH LIGHT WIND FROM WEST, UNSTABLE. 2000 Hr.

DATA VALUE EXTREMES ARE -0.95 2.52

ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL
(MAXIMUM INCLUDED IN HIGHEST LEVEL ONLY)

MINIMUM	BELOW	0.01	0.17	0.32	0.63	1.26	2.50
MAXIMUM	0.01	0.17	0.32	0.63	1.26	2.50	5.00

PERCENTAGE OF TOTAL ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL

3.12	6.12	6.25	12.50	25.00	50.00

FREQUENCY DISTRIBUTION OF DATA POINT VALUES IN EACH LEVEL

LEVEL	1	2	3	4	5	6
SYMBOLS	-	:	+	X	#	@
FREQ.	405	107	17	4	0	0	1

FIGURE C-12

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Light Winds from West
Improvement in SO_2 through "Social Cost" Dispatching



FIGURE C-13

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Light Winds from West
Density of Exposure to Pollution without Regard tc "Social Cost"



FIGURE C-14

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Light Winds from West
Optimal Dispatching Density of Exposure



FIGURE C-15

ST. LOUIS REGION STUDY
Power Plant Study Area
Dispatching under Light Winds from West
Improvement in Population Exposure through "Social Cost" Dispatching





FIGURE D-1

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Suspended Particulate Concentration — Daily Averages
Background Values Added

a. Annual Geometric Mean b. Concentration of Highest Cumulative Frequency Percentile

c. Oblique View of a. d. Oblique View of b.





FIGURE D-2

KANSAS CITY REGION STUDY
Tracted Area of SMSA
Suspended Particulates — Daily Averages
Annual Geometric Mean
a. Background Values Added at 9 Locations
b. Concentration Exceeding Health Standard





FIGURE D-3

KANSAS CITY REGION STUDY
Tracted Area of SMSA
Suspended Particulates — Daily Averages
Concentration of Highest Cumulative Frequency Percentile
a. Background Values Added at 4 Locations
b. Concentration Exceeding Health Standard





FIGURE D-4

KANSAS CITY REGION STUDY
Convex Area of Air Pollution Monitoring Stations
Suspended Particulates — Daily Averages
Annual Geometric Mean
a. Data from 16 Monitoring Stations
b. Background Values Added at 4 Locations



FIGURE D-5

KANSAS CITY REGION STUDY
Convex Area of Air Pollution Monitoring Stations
Suspended Particulates — Daily Averages
Concentration of Highest Cumulative Frequency Percentile
a. Data from 16 Monitoring Stations
b. Background Values Added at 4 Locations



FIGURE D-6

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Dustfall of Settleable Particulates
Geometric Mean of Monthly Averages
a. Data Measured at 36 Monitoring Stations
b. Background Values Added at 4 Points
c. Oblique View of a. d. Oblique View of b.



FIGURE D-7

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Sulfation Rate — Maximum Monthly Average
a. Data from 35 Stations
b. Background Added
c. Corrected for Missing Observations



FIGURE D-8

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Dustfall of Settleable Particulates
Geometric Mean of Monthly Averages
a. Base Map of Monitoring Stations
b. Data Measured at 16 Monitoring Stations
c. Data Measured at 35 Monitoring Stations



FIGURE D-9

KANSAS CITY REGION STUDY
Dustfall of Settleable Particulates
Geometric Mean of Monthly Averages
a. Data Measured at 35 Monitoring Stations
b. Data Measured at 16 Monitoring Stations
c. Comparison of Surfaces from a. and b.



FIGURE D-10

KANSAS CITY REGION STUDY
Suspended Particulates — Daily Averages
Annual Geometric Mean
a. Measured at 16 Monitoring Stations
b. Regression Residuals on Geometric Mean
of Monthly Dustfall



FIGURE D-11

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Sulfation Rate — Annual Arithmetic Mean
a. Data from 35 Monitoring Stations
b. Background Added at 5 Points-Areas Exceeding Standard
c Detail of b. for Tracted Area

FIGURE D-12

KANSAS CITY REGION STUDY
Tracted Area of SMSA
Sulfation Rate — Background Added at 5 Points
Areas Exceeding Health Standard
a. Annual Arithmetic Mean
b. Maximum Monthly Average
c. Combination of a. and b.







FIGURE D-13

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Suspended Particulate Concentration Exceeding Health Standard
a. Concentration of Highest Cumulative Frequency Percentile
b. Annual Geometric Mean of Daily Averages
c. Comparison of a. and b.



FIGURE D-14
KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Dustfall of Settleable Particulates
a. Annual Arithmetic Mean
b. Maximum Monthly Average
c. 3-month Average Calculated from Monthly Data

FIGURE D-15

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
SO_2 Concentration Calculated from Sulfation Rates
a. Yearly Average
b. 24-hour Average Occurring Once Every 3 Months
c. 1-hour Average Occurring Once Every 4 Days



FIGURE D-16

KANSAS CITY REGION STUDY
Averaging Times According to Larsen
a. SO_2 Concentration Calculated from Monthly Sulfation Rates
24-hour Average Occurring Once Every 3 Months
b. SO_2 Concentration Calculated from Monthly Sulfation Rates
1-hour Average Occurring Once Every 4 Days
c. Suspended Particulate Concentration
1-hour Average Occurring Once a Year



FIGURE D-17

KANSAS CITY REGION STUDY
Tracted Area of SMSA
Demographic Data
a. Population Density — Equal Data Points per Level.
b. Population Density — Equal Interval Value Range
c. Median Income



FIGURE D-18

KANSAS CITY REGION STUDY
Tracted Area of SMSA
Dustfall Concentration
Highest 3-month Average Calculated with the Larsen Model
a. Background Values Added at 4 Points
b. Residential and Industrial Areas Exceeding Standards
c. Same as b. Extended over Entire SMSA



FIGURE D-19

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Air Pollutant Emission Density — Sulfur Oxides
a. Average Space Heating Day
b. Maximum Space Heating Day



FIGURE D-20

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Air Pollutant Emission Density
a. Carbon Monoxide — Average Day
b. Particulates — Maximum Space Heating Day

FIGURE D-21

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Particulate Emission Density
a. Average Space Heating Day
b. Maximum Space Heating Day
c. Day without Space Heating





FIGURE D-22

KANSAS CITY REGION STUDY
Tracted Area of SMSA
a. Correlation between Different Air Quality Standards
b. Correlation between Different Social Problem Areas





FIGURE D-23

KANSAS CITY REGION STUDY
Tracted Area of SMSA
Correlation of Air Quality and Demographic Data

a. Levels 1 & 3 — more than 25% of Population Negroes
   Levels 2 & 3 — Any of Air Quality Standards Exceeded
b. Levels 1 & 3 — Annual Geometric Mean Exceeds 75 mg/m^3
   Levels 2 & 3 — Median Age of Population above 35 yrs

FIGURE D-24

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Suspended Particulates — Annual Geometric Mean of Daily Averages
Maximum Search Radius for Interpolation:
a. 1.0 inches b. 1.5 inches c. 2.0 inches



FIGURE D-25

KANSAS CITY REGION STUDY
Standard Metropolitan Statistical Area
Suspended Particulates — Annual Geometric Mean of Daily Averages
Maximum Search Radius for Interpolation:
a. 2.5 inches b. 3.0 inches c. 3.5 inches



FIGURE E-1

MONTREAL REGION STUDY
Distribution of Land Use



FIGURE E-2

MONTREAL REGION STUDY
Index of Atmospheric Purity
Equal Value Range



FIGURE E-3

MONTREAL REGION STUDY
Index of Atmospheric Purity
Equal Distribution of Data Points to Contour Levels



FIGURE E-4

MONTREAL REGION STUDY
Number of Epiphytes as an Air Pollution Index
Equal Value Range



FIGURE E-5

MONTREAL REGION STUDY
Distribution of I.A.P. over the Urbanized Area
Equal Value Range



FIGURE E-6

MONTREAL REGION STUDY
Distribution of I. A. P. over the Urbanized Area
Equal Distribution of Data Points to Contour Lines



FIGURE E-7

MONTREAL REGION STUDY
Coefficient of Haze
City of Montreal Air Pollution Network



FIGURE E-8

MONTREAL REGION STUDY
Sulfur Dioxide
City of Montreal Air Pollution Network



FIGURE E-9

MONTREAL REGION STUDY
Insoluble Residue
City of Montreal Air Pollution Network



FIGURE E-10

MONTREAL REGION STUDY
"Best Selection" Procedure
Distribution of Physcia Millegrana by Coverage & Fertility



FIGURE E-11

MONTREAL REGION STUDY
"Best Selection" Procedure
Distribution of Physcia Millegrana, Physcia Stellaris,
Xanthoria Fallax, & Leskea Polycarpa by Coverage and Fertility



FIGURE E-12

MONTREAL REGION STUDY
Difference between Figures E-11 and E-2
Discrepancies in "Best Selection"



FIGURE E-13

MONTREAL REGION STUDY
"Alternate Selection" Procedure
Distribution of Physcia Millegrana, Physcia Adcendens,
Parmelia Subaurifera, & Xanthoria Fallax by Coverage only



FIGURE E-14

MONTREAL REGION STUDY
Difference between Figures E-13 and E-2
Discrepancies in "Alternate Selection"



FIGURE E-15

MONTREAL REGION STUDY
"Alternate Selection" Procedure by Coverage
Composite of Separate Urban and Rural Mappings



FIGURE F-1

CONNECTICUT STUDY
Connecticut Air Sampling Network
Measured Suspended Particulates Concentration
a. 1966 — 12 months b. 1968 — first 6 months



SYMAP

C CONNECTICUT AIR SAMPLING NETWORK
C SUSPENDED PARTICULATES (UG/M3)
C 1966

DATA VALUE EXTREMES ARE 43.00 110.00

SOURCE- CONNECTICUT STATE DEPARTMENT OF HEALTH
ENVIRONMENTAL HEALTH SERVICES DIVISION
AVERAGE OF DAILY LEVELS

ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL
('MAXIMUM' INCLUDED IN HIGHEST LEVEL ONLY)

MINIMUM	43.00	56.40	69.80	83.20	96.60
MAXIMUM	56.40	69.80	83.20	96.60	110.00

PERCENTAGE OF TOTAL ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL

20.00 20.00 20.00 20.00 20.00

FREQUENCY DISTRIBUTION OF DATA POINT VALUES IN EACH LEVEL

LEVEL	1	2	3	4	5
SYMBOLS					
FREQ.	6	5	4	3	2



SYMAP

C CONNECTICUT AIR SAMPLING NETWORK
C SUSPENDED PARTICULATES (UG/M3)
C 1967

DATA VALUE EXTREMES ARE 49.00 91.00

SOURCE- CONNECTICUT STATE DEPARTMENT OF HEALTH
ENVIRONMENTAL HEALTH SERVICES DIVISION
AVERAGE OF DAILY LEVELS

ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL
('MAXIMUM' INCLUDED IN HIGHEST LEVEL ONLY)

MINIMUM	49.00	57.40	65.80	74.20	82.60
MAXIMUM	57.40	65.80	74.20	82.60	91.00

PERCENTAGE OF TOTAL ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL

20.00 20.00 20.00 20.00 20.00

FREQUENCY DISTRIBUTION OF DATA POINT VALUES IN EACH LEVEL

LEVEL	1	2	3	4	5
SYMBOLS					
FREQ.	9	2	3	4	6



SYMAP

C CONNECTICUT AIR SAMPLING NETWORK
C SUSPENDED PARTICULATES (UG/M3)
C PERCENT CHANGE 1966 - 1967

DATA VALUE EXTREMES ARE -28.00 78.00

SOURCE- CONNECTICUT STATE DEPARTMENT OF HEALTH
ENVIRONMENTAL HEALTH SERVICES DIVISION
AVERAGE OF DAILY LEVELS

ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL
('MAXIMUM' INCLUDED IN HIGHEST LEVEL ONLY)

MINIMUM	-28.00	-10.00	10.00	30.00	50.00
MAXIMUM	-10.00	10.00	30.00	50.00	78.00

PERCENTAGE OF TOTAL ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL

20.00 20.00 20.00 20.00 20.00

FREQUENCY DISTRIBUTION OF DATA POINT VALUES IN EACH LEVEL

LEVEL	1	2	3	4	5
SYMBOLS					
FREQ.	7	3	4	5	4



SYMAP

C PERCENT INCREASE IN SUSPENDED PARTICULATES, 1966-1967
C L = LOW (BELOW ZERO) H = HIGH (ABOVE 50 PERCENT)
C 1 = -15 PERCENT 2 = 15-35 PERCENT 3 = 35-50 PERCENT

DATA VALUE EXTREMES ARE -28.00 78.00

SOURCE- CONNECTICUT STATE DEPARTMENT OF HEALTH
ENVIRONMENTAL HEALTH SERVICES DIVISION
AVERAGE OF DAILY LEVELS

ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL
('MAXIMUM' INCLUDED IN HIGHEST LEVEL ONLY)

MINIMUM	BELOW 0.00	0.00	15.00	35.00	ABOVE 50.00
MAXIMUM	0.00	15.00	35.00	50.00	

PERCENTAGE OF TOTAL ABSOLUTE VALUE RANGE APPLYING TO EACH LEVEL

20.00 40.00 30.00

FREQUENCY DISTRIBUTION OF DATA POINT VALUES IN EACH LEVEL

LEVEL	L	1	2	3	H
SYMBOLS					
FREQ.	9	5	5	6	4

FIGURE F-2

CONNECTICUT STUDY
Connecticut Air Sampling Network
SYMAP Correspondence Course Examples
Measured Suspended Particulates Concentration

a. 1966 b. 1967
c. Percent Change — 1966 to 1967 d. Percent Change — Contour Levels Specified



FIGURE F-3

CONNECTICUT STUDY
Travelers Research Center Emission Inventory
a. Population-related Emissions
b. Automobile-related Emissions



FIGURE F-4

CONNECTICUT STUDY
Travelers Research Center SOx Modeling
a. Down Valley Stable Flow — October Noon
b. Northwest Wind — January Midnight



FIGURE F-5

CONNECTICUT STUDY
Travelers Research Center SOx Modeling
a. Calculated Values — October 15, 1968
b. Observed Values — October 15, 1968





FIGURE F-6

CONNECTICUT STUDY
Travelers Research Center SOx Modeling
Values for October 15, 1968
a. Calculated Greater than Observed
b. Observed Greater than Calculated



FIGURE G-1

SOUTHERN NEW ENGLAND STUDY
Concentration of Suspended Particulates
Based on 206 Selected Urban and Rural Recordings
a. Equal Number of Data Points in Each Level
b. Equal Interval Contour Levels



FIGURE G-2

SOUTHERN NEW ENGLAND STUDY
United States Income Potential
Based on 206 Urban and Rural County Values
a. Equal Number of Data Points in Each Level
b. Equal Interval Contour Levels



FIGURE G-3

SOUTHERN NEW ENGLAND STUDY
Concentration of Suspended Particulates
a. Base Map of Monitoring Stations
and Background Data Points
b. Based on Actual and Estimated Values



FIGURE G-4

SOUTHERN NEW ENGLAND STUDY
Income Potential Calculated at City & County Data Points
a. Surface Based on Counties Only
b. Surface Based on Cities and Counties



FIGURE G-5

SOUTHERN NEW ENGLAND STUDY
Concentration of Suspended Particulates
a. Based on Measured 1966 Values Only
b. Based on Actual and Estimated Values







FIGURE G-6

SOUTHERN NEW ENGLAND STUDY
Income Potential Surface
a. Calculated at City and County Data Points
b. Oblique View of a.
c. Calculated on an 8-mile Grid
d. Oblique View of b.



FIGURE G-7

SOUTHERN NEW ENGLAND STUDY
Comparison of Income Surfaces — Oblique Views
a. Income Density
b. Income Potential at Data Points
c. Income Potential on an 8-mile Grid



FIGURE G-8

SOUTHERN NEW ENGLAND STUDY
Comparison of Population Surfaces
a. Population Density
b. Population Potential at Data Points
c. Population Potential on an 8-mile Grid





FIGURE G-9

SOUTHERN NEW ENGLAND STUDY
Comparison of Air Pollution and Income Potential Surfaces
Oblique Views
a. Concentration of Suspended Particulates
Actual and Estimated Values Used
b. Income Potential — Calculated on an 8-mile Grid







FIGURE G-10

SOUTHERN NEW ENGLAND STUDY
Critical Features of Income Potential Surface
a. Income Potential Surface
b. Hills and Course Lines
c. Dales and Ridge Lines



FIGURE H-1

PUGET SOUND REGION STUDY
Carbon Monoxide Motor Vehicle Emissions
Cities and Corridors Plan — System 35



FIGURE H-2

PUGET SOUND REGION STUDY
Carbon Dioxide Motor Vehicle Emissions
Cities and Corridors Plan — System 35

FIGURE H-3

PUGET SOUND REGION STUDY
Hydrocarbon Motor Vehicle Emissions
Cities and Corridors Plan — System 35



FIGURE H-4

PUGET SOUND REGION STUDY
Carbon Monoxide Motor Vehicle Emissions
Continuation of Existing Trends — System 32



FIGURE H-5

PUGET SOUND REGION STUDY
Differences in Carbon Monoxide Motor Vehicle Emissions
System 35 Exceeds System 32



FIGURE H-6

PUGET SOUND REGION STUDY
Differences in Carbon Monoxide Motor Vehicle Emissions
System 32 Exceeds System 35



FIGURE I-1

LOS ANGELES COUNTY STUDY
Maximum Carbon Monoxide Concentration
Calculated Using 8-hour Values 1956-1967
a. Best Use of Barriers and Mountain Background Points
b. Measured Data Points Only

FIGURE I-2

LOS ANGELES COUNTY STUDY
Maximum Carbon Monoxide Concentration
Calculated Using 8-hour Values 1956-1967
Use of Mountain Background Points and Barrier
a. 2 Full-strenth Barriers and 5 Mountain Points
b. 8 Mountain Points



FIGURE I-3

LOS ANGELES COUNTY STUDY
Maximum Carbon Monoxide Concentration
Calculated Using 8-hour Values 1956-1967
a. Half-strength Barrier on Left; Full-strength on Right
b. Full-strength Barrier; Erroneous Influx from East



FIGURE I-4

LOS ANGELES COUNTY STUDY
Maximum Carbon Monoxide Concentration
Calculated Using 8-hour Values 1956-1967
a. 2 Full-strength Barriers
b. Modeling Flow through Mountain Pass



FIGURE I-5

LOS ANGELES COUNTY STUDY
Maximum Carbon Monoxide Concentration
Calculated Using 8-hour Values
a. 1956-1967 b. 1958



FIGURE I-6

LOS ANGELES COUNTY STUDY
Maximum Carbon Monoxide Concentration
Calculated Using 8-hour Values
a. 1959 b. 1960



FIGURE I-7

LOS ANGELES COUNTY STUDY
Maximum Carbon Monoxide Concentration
Calculated Using 8-hour Values
a. 1961 b. 1962



FIGURE I-8

LOS ANGELES COUNTY STUDY
Carbon Monoxide Concentration
Calculated Using 8-hour Values 1956-1967
a. Geometric Mean
b. Percent of Time Concentration Exceeds 0.1 p.p.m.

FIGURE I-9

SAN BERNARDINO COUNTY STUDY
Total Annual Suspended Dust Concentration
1966



FIGURE I-10

SAN BERNARDINO COUNTY STUDY
Total Annual Suspended Dust Concentration
a. 1966 b. 1967
c. Absolute Increase, 1966 to 1967



FIGURE I-11

SAN BERNARDINO COUNTY STUDY
Total Annual Suspended Dust Concentration
a. Spring b. Spring c. Fall



FIGURE I-12

CALIFORNIA STUDY
Maximum Hourly Average Concentration of Oxidant
a. November 1967 b. December 1967 c. January 1968



FIGURE I-13

CALIFORNIA STUDY
Maximum Hourly Average Concentration of Oxidant
a. February 1968 b. March 1968 c. April 1968



FIGURE I-14

CALIFORNIA STUDY
Maximum Hourly Average Concentration of Oxidant
a. May 1968 b. June 1968 c. July 1968



FIGURE J-1

BOSTON REGION STUDY
1966 Concentration of Suspended Particulates
a. January b. February c. March



FIGURE J-2
BOSTON REGION STUDY
1966 Concentration of Suspended Particulates
a. April b. May c. June



FIGURE J-3

BOSTON REGION STUDY
1966 Concentration of Suspended Particulates
a. July b. August c. September



FIGURE J-4

BOSTON REGION STUDY
1966 Concentration of Suspended Particulates
a. October b. November
c. 11-month Average



FIGURE J-5

BOSTON REGION STUDY
1966 Concentration of SO_2
a. January b. February
c. Average of a. and b. (Winter)



FIGURE J-6

BOSTON REGION STUDY
1966 Concentration of SO_2
a. April b. May
c. Average of a. and b. (Spring)



FIGURE J-7

BOSTON REGION STUDY
1966 Concentration of SO_2
a. July b. August
c. Average of a. and b. (Summer)



FIGURE J-8

BOSTON REGION STUDY
1966 Concentration of SO_2
a. October b. November
c. Average of a. and b. (Fall)



FIGURE J-9

BOSTON REGION STUDY
1966 Concentration of SO_2
a. Episode Period — November 21, 1966
b. Episode Period — November 24, 1966
c. January-November (11-month) Average



FIGURE J-10

BOSTON REGION STUDY
Development of Emission Inventory
a. Population Density by Towns
b. SOx Emissions, Average Day
c. Suspended Particulate Emissions — Average Day



FIGURE X-1

PHILADELPHIA AIR QUALITY CONTROL REGION
Calculated SOx — 12-hour Half-life
Oblique Views 1 and 2 of Surface



FIGURE X-2

PHILADELPHIA AIR QUALITY CONTROL REGION
Calculated SOx — 12-hour Half-life
Oblique Views 3 and 4 of Surface